arXiv:1610.02367v1 [math-ph] 7 Oct 2016

Matrix commuting differential operators of rank 2

Vardan Oganesyan

Abstract

In this paper we propose a very effective method for constructing matrix commuting differential operators of rank 2 and vector rank (2,2). We find new matrix commuting differential operators L, M of orders 2 and $2g$ respectively.

Introduction

Let us consider two differential operators

$$L_n = \sum_{i=0}^{n} u_i(x)\partial_x^i, \quad L_m = \sum_{i=0}^{m} v_i(x)\partial_x^i,$$

where coefficients $u_i(x)$ and $v_i(x)$ are scalar or matrix valued functions. The commutativity condition $L_nL_m = L_mL_n$ is equivalent to a very complicated system of nonlinear differential equations. The theory of commuting ordinary differential operators was first developed in the beginning of the XX century in the works of Wallenberg [1], Schur [2].

If two differential operators with scalar or matrix valued coefficients commute, then there exists a nonzero polynomial $R(z, w)$ such that $R(L_n, L_m) = 0$ (see [3], [4]). The curve Γ defined by $R(z, w) = 0$ is called the *spectral curve*. If

$$L_n\psi = z\psi, \quad L_m\psi = w\psi,$$

then $(z, w) \in \Gamma$.

If coefficients are scalar functions, then for almost all $(z, w) \in \Gamma$, the dimension of the space of common eigenfunctions ψ is the same. The dimension of the space of common eigenfunctions of two commuting scalar differential operators is called the *rank* of this pair. The rank is a common divisor of m and n. The genus of the spectral curve of a pair of commuting operators is called the genus of this pair.

This research was supported by the Russian Science Foundation under grant 16-11-10260 and was done at the Faculty of Mechanics and Mathematics, Department of Geometry and Topology of Lomonosov Moscow State University

If the rank of two commuting scalar differential operators equals 1, then there are explicit formulas for coefficients of commutative operators in terms of Riemann theta-functions (see [5]).

The case when rank of scalar commuting operators is greater than 1 is much more difficult. The first examples of commuting scalar differential operators of the nontrivial rank 2 and the nontrivial genus $g = 1$ were constructed by Dixmier [6] for the nonsingular elliptic spectral curve.

A general classification of commuting scalar differential operators was obtained by Krichever [7]. The general form of commuting scalar operators of rank 2 for an arbitrary elliptic spectral curve was found by Krichever and Novikov [8]. The general form of scalar commuting operators of rank 3 with arbitrary elliptic spectral curve was found by Mokhov [9],[10]. In [11] Mironov developed theory of self-adjoint scalar operators of rank 2 and found examples of commuting scalar operators of rank 2 and arbitrary genus. Using Mironov's method many examples of scalar commuting operators of rank 2 and arbitrary genus were found (see [12], [13], [14], [15], [16]). Moreover, examples of commuting scalar differential operators of arbitrary genus and arbitrary rank with polynomial coefficients were constructed by Mokhov in [17], [18].

Theory of commuting differential operators helps to find solutions of nonlinear partial differential equations from mathematical physics (see [19], [20], [21], [22]). Also there are deep connections between theory of commuting scalar differential operators and Schottky problem(see [23], [24]). The theory of commuting differential operators with polynomial coefficients has connections with the Dixmier conjecture and Jacobian conjecture (see [25], [26]).

A general classification of commuting matrix differential operators was obtained by Grinevich [4]. Grinevich considered two differential operators

$$L = \sum_{i=0}^{m} U_i \partial_x^i, \quad M = \sum_{i=0}^{n} V_i \partial_x^i,$$

where U_i and V_i are smooth and complex-valued $s \times s$ matrices. Let us suppose the following conditions

1) $det(U_m) \neq 0$.
2) Eigenvalues $\lambda_1(x), ..., \lambda_s(x)$ of U_m are distinct.
3) Matrix V_n is diagonalizable. Let $\mu_1(x), ..., \mu_s(x)$ be eigenvalues of matrix V_n. Suppose that functions $\dfrac{\mu_i^m}{\lambda_i^n}$ are distinct constants for all $i = 1, ..., s$.

Easy to see that if L and M commute, then U_m and V_n commute. If operators

L and M commute, then FLF^{-1} and FMF^{-1} commute, where F is matrix. We also can change variable. So, without loss of generality we can suppose that

$$(U_m)_{ij} = \delta_{ij}\lambda_i, \quad (V_n)_{ij} = \delta_{ij}\mu_i, \quad tr(U_{m-1}) = 0$$

Let Γ be the spectral curve of commuting matrix operators L, M. Spectral curve of matrix commuting operators can be reducible. Let Γ_i be an irreducible component of the spectral curve. The dimension of the space of common eigenfunctions

$$L\psi = z\psi, \quad M\psi = w\psi, \quad (z, w) \in \Gamma_i$$

is called the rank of commuting pair on Γ_i. Grinevich discovered that the spectral curve Γ has s points at infinity. So, $\Gamma = \bigcup_{i=1}^{k} \Gamma_i$, where $k \leqslant s$. Let l_i be the rank of operators on Γ_i. Operators L, M are called commuting operators of vector rank $(l_1, ..., l_k)$, where $k \leqslant s$. Numbers l_i are common divisors of m and n. For more details see [4]. Also see [27], [28], [29].

If the rank of commuting matrix differential operators equals 1, then there exists explicit formulas for coefficients in terms of Riemann theta-functions [30].

In this paper we propose a very effective method for constructing matrix commuting operators of rank 2 and vector rank (2,2). We find new commuting operators L, M of orders 2 and $2g$ respectively.

Acknowledgments

The author wishes to express gratitude to Professor O. I. Mokhov for advices and help in writing this paper.

Explicit examples of commuting matrix differential operators of rank 2

Let us consider the operator

$$L = E(x)\partial_x^2 + R(x)\partial + Q(x), \tag{1}$$

where

$$E = \begin{pmatrix} \lambda_1(x) & \lambda_3(x) \\ 0 & \lambda_2(x) \end{pmatrix}, \quad R = \begin{pmatrix} r_1(x) & r_3(x) \\ r_2(x) & -r_1(x) \end{pmatrix}, \quad Q = \begin{pmatrix} q_1(x) & q_3(x) \\ q_2(x) & q_4(x) \end{pmatrix}$$

We want to find an operator M of order $2g$ such that $[L, M] = 0$. Let us consider the operator

$$M = B_0(x)L^g + (A_1(x)\partial_x + B_1(x))L^{g-1} + (A_2(x)\partial_x + B_2(x))L^{g-2} + ...+ \\ +(A_{g-1}(x)\partial_x + B_{g-1}(x))L + A_g(x)\partial_x + B_g(x), \tag{2}$$

where

$$A_{g-k} = \begin{pmatrix} a_1^{g-k}(x) & a_3^{g-k}(x) \\ a_2^{g-k}(x) & a_4^{g-k}(x) \end{pmatrix}, \quad B_{g-k} = \begin{pmatrix} b_1^{g-k}(x) & b_3^{g-k}(x) \\ b_2^{g-k}(x) & b_4^{g-k}(x) \end{pmatrix}, \quad A_0 = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}.$$

We note that in the formulas above the number $g - k$ is **index not degree**. The index $g - k$ says us that functions a_i^{g-k} and b_i^{g-k} are elements of matrices A_{g-k} and B_{g-k} respectively.

Let us note that if $[L, N] = 0$, then $[L, MN] = LMN - MNL = LMN - MLN = [L, M]N$, where L, M, N are matrix differential operators. So, we see that

$$[L, M] = [L, \sum_{k=0}^{g}(A_{g-k}\partial_x + B_{g-k})L^k] = \sum_{k=0}^{g}[L, (A_{g-k}\partial_x + B_{g-k})L^k] = \\ = \sum_{k=0}^{g}[L, (A_{g-k}\partial_x + B_{g-k})]L^k.$$

Direct calculations show that

$$\big[L, \ \ A_{g-k}\partial_x + B_{g-k}\big] = [E\partial_x^2 + R\partial + Q, \ \ A_{g-k}\partial_x + B_{g-k}\big] =$$

$$+(EA_{g-k} - A_{g-k}E)\partial_x^3 + (2EA'_{g-k} + EB_{g-k} + RA_{g-k} - A_{g-k}E' - A_{g-k}R - B_{g-k}E)\partial_x^2 + \\ +(EA''_{g-k} + 2EB'_{g-k} + R_{g-k}A' + RB_{g-k} + QA_{g-k} - A_{g-k}R' - A_{g-k}Q - B_{g-k}R)\partial_x + \\ +(EB''_{g-k} + RB'_{g-k} + QB_{g-k} - A_{g-k}Q' - B_{g-k}Q) =$$

$$= K_{g-k}\partial_x^3 + P_{g-k}\partial_x^2 + T_{g-k}\partial_x + F_{g-k},$$

where

$$K_{g-k} = EA_{g-k} - A_{g-k}E,$$
$$P_{g-k} = 2EA'_{g-k} + EB_{g-k} + RA_{g-k} - A_{g-k}E' - A_{g-k}R - B_{g-k}E,$$
$$T_{g-k} = EA''_{g-k} + 2EB'_{g-k} + R_{g-k}A' + RB_{g-k} + QA_{g-k} - A_{g-k}R' - A_{g-k}Q - B_{g-k}R,$$
$$F_{g-k} = EB''_{g-k} + RB'_{g-k} + QB_{g-k} - A_{g-k}Q' - B_{g-k}Q.$$

Using the fact that $\partial_x L = E\partial_x^3 + (E' + R)\partial_x^2 + (R' + Q)\partial_x + Q'$ we get

$$K_{g-k}\partial_x^3 + P_{g-k}\partial_x^2 + T_{g-k}\partial_x + F_{g-k} =$$

$$K_{g-k}E^{-1}\partial_x L + \left(P_{g-k} - K_{g-k}E^{-1}(E' + R)\right)\partial_x^2 +$$
$$\left(T_{g-k} - K_{g-k}E^{-1}(R' + Q)\right)\partial_x + (F_{g-k} - K_{g-k}E^{-1}Q') =$$

$$K_{g-k}E^{-1}\partial_x L + \left(P_{g-k} - K_{g-k}E^{-1}(E' + R)\right)E^{-1}L +$$
$$\left(T_{g-k} - K_{g-k}E^{-1}(R' + Q) - \left(P_{g-k} - K_{g-k}E^{-1}(E' + R)\right)E^{-1}R\right)\partial_x + +$$
$$\left(F_{g-l} - K_{g-k}E^{-1}Q' - \left(P_{g-k} - K_{g-k}E^{-1}(E' + R)\right)E^{-1}Q\right) =$$

$$\widetilde{K}_{g-k}\partial_x L + \widetilde{P}_{g-k}L + \widetilde{T}_{g-k}\partial_x + \widetilde{F}_{g-k},$$

where

$$\begin{gathered}
\widetilde{K}_{g-k} = K_{g-k}E^{-1} \\
\widetilde{P}_{g-k} = \left(P_{g-k} - K_{g-k}E^{-1}(E' + R)\right)E^{-1} \\
\widetilde{T}_{g-k} = \left(T_{g-k} - K_{g-k}E^{-1}(R' + Q) - K_{g-k}E^{-1}(E' + R)E^{-1}R\right) \\
\widetilde{F}_{g-k} = \left(F_{g-k} - K_{g-k}E^{-1}Q' - K_{g-k}E^{-1}(E' + R)E^{-1}Q\right).
\end{gathered} \tag{3}$$

Finally we obtain

$$[L, M] = \sum_{k=0}^{g}[L, (A_{g-k}\partial_x + B_{g-k})]L^k =$$

$$= \left(\widetilde{K}_0\partial_x + \widetilde{P}_0\right)L^{g+1} + \left((\widetilde{T}_0 + \widetilde{K}_1)\partial_x + (\widetilde{F}_0 + \widetilde{P}_1)\right)L^{g-1} +$$
$$+ \left((\widetilde{T}_1 + \widetilde{K}_2)\partial_x + (\widetilde{F}_1 + \widetilde{P}_2)\right)L^{g-1} + ... +$$
$$+ \left((\widetilde{T}_{g-1} + \widetilde{K}_g)\partial_x + (\widetilde{F}_{g-1} + \widetilde{P}_g)\right)L + T_g\partial_x + F_g.$$

So, if

$$\begin{gathered}
\widetilde{K}_0 = 0, \widetilde{P}_0 = 0, \widetilde{K}_1 = -\widetilde{T}_0, \widetilde{P}_1 = -\widetilde{F}_0, ..., \\
\widetilde{K}_m = -\widetilde{T}_{m-1}, \widetilde{P}_m = -\widetilde{F}_{m-1}, ..., \\
\widetilde{K}_g = -\widetilde{T}_{g-1}, \widetilde{P}_g = -\widetilde{F}_{g-1}, \\
\widetilde{T}_g = 0, \widetilde{F}_g = 0,
\end{gathered} \tag{4}$$

then $[L, M] = 0$.

Let us calculate $\widetilde{K}_{g-k}, \widetilde{P}_{g-k}, \widetilde{T}_{g-k}, \widetilde{F}_{g-k}$ from (3). This formulas are too hard for analyzing but in the sequel only special cases are considered. We are going to show that in some cases formulas (4) give a very effective methods for finding commuting operators. Let us describe the main idea. We know that $A_0 = 0$ because M is operator of order $2g$. Let us take B_0 such that $\widetilde{K}_0 = 0$ and $\widetilde{P}_0 = 0$. Then we can calculate $\widetilde{T}_0, \widetilde{F}_0$. Using (4) we can find $\widetilde{K}_1, \widetilde{P}_1$, then $a_2^1, a_3^1, a_4^1, b_1^1, b_2^1, b_3^1, b_4^1$. And using (4) we can find $\widetilde{T}_1$ and $\widetilde{F}_1$. So, we get recurrence relations

$$\begin{cases} a_i^{m+1} = g_i(a_1^m, a_2^m, a_3^m, a_4^m, b_1^m, b_2^m, b_3^m, b_4^m, r_1, r_2, r_3, q_1, q_2, q_3, q_4), & i = 1, 2, 3, 4 \\ b_i^{m+1} = h_i(a_1^m, a_2^m, a_3^m, a_4^m, b_1^m, b_2^m, b_3^m, b_4^m, r_1, r_2, r_3, q_1, q_2, q_3, q_4) & i = 1, 2, 3, 4. \end{cases}$$

We will see that if there exists g such that

$$\begin{cases} a_i^{g+1} = 0 \\ b_i^{g+1} = 0 \end{cases} \quad i = 1, ..., 4$$

then the operator L commutes with operator M.

Let us suppose that $\lambda_1 = 1$, $\lambda_2 = -1$, $\lambda_3 = 0$, $r_2(x) = r_3(x) = 0$, $q_3 = q_2$ and $q_4 = -q_1$. Then we have

$$\widetilde{K}_{g-k} = \begin{pmatrix} 0 & -2a_3^{g-k} \\ -2a_2^{g-k} & 0 \end{pmatrix}, \quad \widetilde{P}_{g-k} = \begin{pmatrix} 2(a_1^{g-k})' & -2b_3^{g-k} - 2(a_3^{g-k})' \\ -2b_2^{g-k} - 2(a_2^{g-k})' & 2(a_4^{g-k})' \end{pmatrix},$$

$$\widetilde{T}_{g-k} = \begin{pmatrix} \widetilde{T}^1_{g-k,1} & \widetilde{T}^1_{g-k,2} \\ \widetilde{T}^2_{g-k,1} & \widetilde{T}^2_{g-k,2} \end{pmatrix}, \quad \widetilde{F}_{g-k} = \begin{pmatrix} \widetilde{F}^1_{g-k,1} & \widetilde{F}^1_{g-k,2} \\ \widetilde{F}^2_{g-k,1} & \widetilde{F}^2_{g-k,2} \end{pmatrix},$$

$$\begin{aligned} \widetilde{T}^1_{g-k,1} &= a_2^{g-k}q_2 + a_3^{g-k}q_2 - r_1(a_1^{g-k})' + 2(b_1^{g-k})' - a_1^{g-k}r_1' + (a_1^{g-k})'', \\ \widetilde{T}^1_{g-k,2} &= -a_1^{g-k}q_2 + a_4^{g-k}q_2 - r_1(a_3^{g-k})' + 2(b_3^{g-k})' - a_3^{g-k}r_1' + (a_3^{g-k})'', \\ \widetilde{T}^2_{g-k,1} &= a_1^{g-k}q_2 - a_4^{g-k}q_2 + r_1(a_2^{g-k})' - 2(b_2^{g-k})' + a_2^{g-k}r_1' - (a_2^{g-k})'', \\ \widetilde{T}^2_{g-k,2} &= a_2^{g-k}q_2 + a_3^{g-k}q_2 + r_1(a_4^{g-k})' - 2(b_4^{g-k})' + a_4^{g-k}r_1' - (a_4^{g-k})'', \end{aligned}$$

$$\begin{aligned} \widetilde{F}^1_{g-k,1} &= b_2^{g-k}q_2 + b_3^{g-k}q_2 - 2q_1(a_1^{g-k})' + 2q_2(a_3^{g-k})' + r_1(b_1^{g-k})' - q_1'a_1^{g-k} + q_2'a_3^{g-k} + (b_1^{g-k})'', \\ \widetilde{F}^1_{g-k,2} &= -b_1^{g-k}q_2 + b_4^{g-k}q_2 - 2q_2(a_1^{g-k})' - 2q_1(a_3^{g-k})' + r_1(b_3^{g-k})' - q_1'a_3^{g-k} - q_2'a_1^{g-k} + (b_3^{g-k})'', \\ \widetilde{F}^2_{g-k,1} &= b_1^{g-k}q_2 - b_4^{g-k}q_2 + 2q_1(a_2^{g-k})' - 2q_2(a_4^{g-k})' - r_1(b_2^{g-k})' + q_1'a_2^{g-k} - q_2'a_4^{g-k} - (b_2^{g-k})'', \\ \widetilde{F}^2_{g-k,2} &= b_2^{g-k}q_2 + b_3^{g-k}q_2 + 2q_2(a_2^{g-k})' + 2q_1(a_4^{g-k})' - r_1(b_4^{g-k})' + q_1'a_4^{g-k} + q_2'a_2^{g-k} - (b_4^{g-k})'' \end{aligned}$$

From (4) we get

$$\widetilde{K}_{g-k+1} = -\widetilde{T}_{g-k}, \widetilde{P}_{g-k+1} = -\widetilde{F}_{g-k}.$$

So, we obtain recurrence relations

$$\begin{gathered} a_i^0(x) \equiv 0, \qquad i = 1, ...4, \\ b_2^0(x) = b_3^0 \equiv 0 \end{gathered} \tag{5}$$

$$\begin{gathered} \widetilde{T}^1_{g-k,1} = 0 \Leftrightarrow \\ b_1^{g-k} = -\frac{1}{2}\int \left(a_2^{g-k}q_2 + a_3^{g-k}q_2 - (a_1^{g-k})'r_1 - a_1^{g-k}r_1' + (a_1^{g-k})''\right)dx + C_1^{g-k}, \end{gathered} \tag{6}$$

$$\begin{gathered} \widetilde{T}^2_{g-k,2} = 0 \Leftrightarrow \\ b_4^{g-k} = \frac{1}{2}\int \left(a_2^{g-k}q_2 + a_3^{g-k}q_2 + (a_4^{g-k})'r_1 + a_4^{g-k}r_1' - (a_4^{g-k})''\right)dx + C_2^{g-k}, \end{gathered} \tag{7}$$

$$\begin{gathered} -2a_3^{g-k+1} = -\widetilde{T}^1_{g-k,2} \Leftrightarrow \\ a_3^{g-k+1} = \frac{1}{2}\left(-a_1^{g-k}q_2 + a_4^{g-k}q_2 - (a_3^{g-k})'r_1 + 2(b_3^{g-k})' - a_3^{g-k}r_1' + (a_3^{g-k})''\right), \end{gathered} \tag{8}$$

$$\begin{gathered} -2a_2^{g-k+1} = -\widetilde{T}^2_{g-k,1} \Leftrightarrow \\ a_2^{g-k+1} = \frac{1}{2}\left(a_1^{g-k}q_2 - a_4^{g-k}q_2 + (a_2^{g-k})'r_1 - 2(b_2^{g-k})' + a_2^{g-k}r_1' - (a_2^{g-k})''\right), \end{gathered} \tag{9}$$

$$\begin{gathered} 2(a_1^{g-k+1})' = -\widetilde{F}^1_{g-k,1} \Leftrightarrow \\ a_1^{g-k+1} = -\frac{1}{2}\int \left(b_2^{g-k}q_2 + b_3^{g-k}q_2 - 2(a_1^{g-k})'q_1 + 2q_2(a_3^{g-k})' + (b_1^{g-k})'r_1 - \right. \\ \left. -a_1^{g-k}q_1' + a_3^{g-k}q_2' + (b_1^{g-k})''\right)dx + C_3^{g-k+1}, \end{gathered} \tag{10}$$

$$\begin{gathered} 2(a_4^{g-k+1})' = -\widetilde{F}^2_{g-k,2} \Leftrightarrow \\ a_4^{g-k+1} = -\frac{1}{2}\int \left(b_2^{g-k}q_2 + b_3^{g-k}q_2 + 2(a_2^{g-k})'q_2 + 2(a_4^{g-k})'q_1 - (b_4^{g-k})'r_1 + \right. \\ \left. +a_4^{g-k}q_1' + a_2^{g-k}q_2' - (b_4^{g-k})''\right)dx + C_4^{g-k+1}, \end{gathered} \tag{11}$$

$$-2b_3^{g-k+1}-(a_3^{g-k+1})'=-\widetilde{F}^1_{g-k,2}\Leftrightarrow$$
$$b_3^{g-k+1}=\frac{1}{2}\Big(-b_1^{g-k}q_2+b_4^{g-k}q_2-2(a_1^{g-k})'q_2-2(a_3^{g-k})'q_1+(b_3^{g-k})'r_1-\\ -a_3^{g-k}q_1'-a_1^{g-k}q_2'+(b_3^{g-k})''\Big)-(a_3^{g-k+1})', \tag{12}$$

$$-2b_2^{g-k+1}-(a_2^{g-k+1})'=-\widetilde{F}^2_{g-k,1}\Leftrightarrow$$
$$b_2^{g-k+1}=\frac{1}{2}\Big(b_1^{g-k}q_2-b_4^{g-k}q_2+2(a_2^{g-k})'q_1-2(a_4^{g-k})'q_2-(b_2^{g-k})'r_1+\\ +a_2^{g-k}q_1'-a_4^{g-k}q_2'-(b_2^{g-k})''\Big)-(a_2^{g-k+1})'. \tag{13}$$

We see that

$$\begin{cases}\widetilde{T}_g=0\\ \widetilde{F}_g=0\end{cases}\quad\Leftrightarrow\quad\begin{cases}\widetilde{K}_{g+1}=0\\ \widetilde{P}_{g+1}=0\end{cases}$$

where

$$\widetilde{K}_{g+1}=\begin{pmatrix}0 & -2a_3^{g+1}\\ 2a_2^{g+1} & 0\end{pmatrix},\quad \widetilde{P}_{g+1}=\begin{pmatrix}2(a_1^{g+1})' & -2b_3^{g+1}-2(a_3^{g+1})'\\ -2b_2^{g+1}-2(a_2^{g+1})' & 2(a_4^{g+1})',\end{pmatrix}$$

where C_i^j are arbitrary constants. Let us note that if $a_i^0(x)=0$, $b_2^0(x)=0, b_3^0(x)=0$ for all $i=1,...,4$, then from (6) and (7) we get that $b_1^0(x)=const$ and $b_4^0(x)=const$.

We obtain the following theorem

Theorem 1. *If there exists number g and constants of integration C_j^m such that $a_i^{g+1}=0, b_2^{g+1}=0, b_3^{g+1}=0$ for all $i=1,...,4$, then the operator*

$$L=\begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix}\partial_x^2+\begin{pmatrix}r_1(x) & 0\\ 0 & -r_1(x)\end{pmatrix}\partial_x+\begin{pmatrix}q_1(x) & q_2(x)\\ q_2(x) & -q_1(x)\end{pmatrix}$$

commutes with operator

$$M=B_0L^g+(A_1\partial_x+B_1)L^{g-1}+...+A_0\partial_x+B_0, \tag{14}$$

$$B_0=\begin{pmatrix}\mu_1 & 0\\ 0 & \mu_2\end{pmatrix},\quad B_1=\begin{pmatrix}C_1^1+\dfrac{C_3^1r_1(x)}{2} & -\dfrac{(\mu_1-\mu_2)q_2}{2}\\ \dfrac{(\mu_1-\mu_2)q_2}{2} & C_1^4+\dfrac{C_4^1r_1(x)}{2}\end{pmatrix},$$

where μ_1, μ_2 are arbitrary constants and C_i^j are some constants. We see that if $\mu_1 \neq \mu_2$ and $q_2 \neq const$, then B_1 is not constant matrix hence M is not polynomial in L.

Theorem 2. *The operator*

$$L = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \partial_x^2 + \begin{pmatrix} \alpha_2 x^2 + \alpha_0 & 0 \\ 0 & -\alpha_2 x^2 - \alpha_0 \end{pmatrix} \partial_x + \begin{pmatrix} \beta x^2 + \alpha_2 x & \gamma x \\ \gamma x & -\beta x^2 - \alpha_2 x \end{pmatrix},$$

where

$$\gamma^2 = -n^2 \alpha_2^2, \quad n \in \mathbb{N}$$

and $\alpha_2, \alpha_0, \beta$ are arbitrary constants, commutes with differential operator (14) of order $4n$, where $g = 2n$. The order of operator M equals $4n$.

Remark. Calculations show that if $n \leqslant 3$, then the spectral curve of operators L, M from Theorem 2 is nonsingular for almost all $\alpha_0, \alpha_2, \beta$ and is hyperelliptic. Hence L and M are operators of rank 2. In some cases spectral curve is reducible and we get commuting operators of rank $(2, 2)$. Note that operators from Theorem 2 can't be operators of rank 1. Also note that from Theorem 1 we see that the matrix operator M from Theorem 2 is operator with polynomial coefficients.

Example 1. If $n = 1$ and $\mu_1 = 1, \mu_2 = -1$, then the operator

$$L = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \partial_x^2 + \begin{pmatrix} \alpha_2 x^2 + \alpha_0 & 0 \\ 0 & -\alpha_2 x^2 - \alpha_0 \end{pmatrix} \partial_x + \begin{pmatrix} \beta x^2 + \alpha_2 x & i\alpha_2 x \\ i\alpha_2 x & -\beta x^2 - \alpha_2 x \end{pmatrix}$$

commutes with operator $M = B_0 L^2 + A_1 \partial_x L + B_1 L + A_2 \partial_x + B_2$. Calculations show that

$$M = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \partial_x^4 + \begin{pmatrix} 2(\alpha_2 x^2 + \alpha_0) & 0 \\ 0 & -2(\alpha_2 x^2 + \alpha_0) \end{pmatrix} \partial_x^3 +$$
$$+ \begin{pmatrix} \alpha_2^2 x^4 + 2(\alpha_0 \alpha_2 + \beta) x^2 + 6\alpha_2 x + \alpha_0^2 & i\alpha_2 x \\ i\alpha_2 x & -\alpha_2^2 x^4 - 2(\alpha_0 \alpha_2 + \beta) x^2 - 6\alpha_2 x - \alpha_0^2 \end{pmatrix} \partial_x^2 +$$
$$+ \begin{pmatrix} m_1 & m_2 \\ m_2 & -m_1 \end{pmatrix} \partial_x + \begin{pmatrix} h_1 & h_2 \\ h_2 & -h_1 + 2\beta - \alpha_0 \alpha_2 \end{pmatrix} + C_1 L + \begin{pmatrix} C_0 & 0 \\ 0 & C_0 \end{pmatrix},$$

$$m_1 = 2\alpha_2 \beta x^4 + 4\alpha_2^2 x^3 + 2\alpha_0 \beta x^2 + 4(\alpha_0 \alpha_2 + \beta) x + 4\alpha_2,$$
$$m_2 = i\alpha_2^2 x^3 + i\alpha_0 \alpha_2 x + i\alpha_2,$$
$$h_1 = \beta_2^2 x^4 + 4\alpha_2 \beta x^3 + \frac{3\alpha_2^2}{2} x^2 + 2\alpha_0 \beta x + 4\beta,$$

$$h_2 = i\alpha_2\beta x^3 + \frac{3}{2}i\alpha_2^2 x^2 + \frac{i\alpha_2\alpha_0}{2},$$

where C_1 and C_0 are arbitrary constants. The spectral curve of operators L, M has the form

$$\left(w - C_1 z - (C_0 - \frac{\alpha_2\alpha_0 - 2\beta}{2})\right)^2 = z^4 - (\alpha_0\alpha_2 - 2\beta)z^2 - \alpha_2\alpha_0\beta + \beta^2.$$

If we take $C_0 = \dfrac{\alpha_2\alpha_0 - 2\beta}{2}$, $C_1 = 0$, then we get

$$w^2 = z^4 - (\alpha_0\alpha_2 - 2\beta)z^2 - \alpha_2\alpha_0\beta + \beta^2.$$

This spectral curve is nonsingular if $\alpha_2\alpha_0\beta(\alpha_2\alpha_0 - \beta) \neq 0$. So in nonsingular case we get that operators L, M are operators of rank 2. If $\alpha_0 = 0$, then the spectral curve has the form

$$w^2 = (z^2 + \beta)^2 \Leftrightarrow (w - z^2 - \beta)(w + z^2 + \beta) = 0 \tag{15}$$

We see that if $\alpha_0 = 0$, then the spectral curve is reducible. Note that $M \neq L^2 + \beta$ and $M \neq -L^2 - \beta$ but $(M - L^2 - \beta)(M + L^2 + \beta) = 0$ and we have operators of vector rank (2,2).

Example 2. If $n = 1$ and $\mu_1 = 1, \mu_2 = 2$, then the operator

$$L = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}\partial_x^2 + \begin{pmatrix} \alpha_2 x^2 + \alpha_0 & 0 \\ 0 & -\alpha_2 x^2 - \alpha_0 \end{pmatrix}\partial_x + \begin{pmatrix} \beta x^2 + \alpha_2 x & i\alpha_2 x \\ i\alpha_2 x & -\beta x^2 - \alpha_2 x \end{pmatrix}$$

commutes with operator $M = B_0 L^2 + A_1\partial_x L + B_1 L + A_2\partial_x + B_2$. Direct calculations show that

$$M = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}\partial_x^4 + \begin{pmatrix} 2(\alpha_2 x^2 + \alpha_0) & 0 \\ 0 & 4(\alpha_2 x^2 + \alpha_0) \end{pmatrix}\partial_x^3 +$$

$$+ \begin{pmatrix} \alpha_2^2 x^4 + 2(\alpha_0\alpha_2 + \beta)x^2 + 6\alpha_2 x + \alpha_0^2 & -\dfrac{i\alpha_2 x}{2} \\ -\dfrac{i\alpha_2 x}{2} & 2\alpha_2^2 x^4 + 4(\alpha_0\alpha_2 + \beta)x^2 + 12\alpha_2 x + 2\alpha_0^2 \end{pmatrix}\partial_x^2 +$$

$$+ \begin{pmatrix} m_1 & m_3 \\ m_2 & m_4 \end{pmatrix}\partial_x + \begin{pmatrix} h_1 & h_3 \\ h_2 & h_4 \end{pmatrix} + C_1 L + \begin{pmatrix} C_0 & 0 \\ 0 & C_0 \end{pmatrix},$$

$$m_1 = 2\alpha_2\beta x^4 + 4\alpha_2^2 x^3 + 2\alpha_0\beta x^2 + 4(\alpha_0\alpha_2 + \beta)x + 4\alpha_2,$$

$$m_2 = -\frac{i\alpha_2^2 x^3}{2} - \frac{i\alpha_0\alpha_2 x}{2} - \frac{7i\alpha_2}{2},$$

$$m_3 = -\frac{i\alpha_2^2 x^3}{2} - \frac{i\alpha_0\alpha_2 x}{2} + \frac{5i\alpha_2}{2},$$

$$m_4 = 4\alpha_2\beta x^4 + 8\alpha_2^2 x^3 + 4\alpha_0\beta x^2 + 8(\alpha_0\alpha_2 + \beta)x + 8\alpha_2,$$
$$h_1 = \beta^2 x^4 + 4\alpha_2\beta x^3 + \frac{3\alpha_2^2}{4}x^2 + 2\alpha_0\beta x + \beta + \frac{3\alpha_2\alpha_0}{2},$$
$$h_2 = -\frac{i\alpha_2\beta x^3}{2} - \frac{9i\alpha_2^2 x^2}{4} - \frac{7i\alpha_2\alpha_0}{4},$$
$$h_3 = -\frac{i\alpha_2\beta x^3}{2} + \frac{3i\alpha_2^2 x^2}{4} + \frac{5i\alpha_2\alpha_0}{4},$$
$$h_4 = 2\beta^2 x^4 + 8\alpha_2\beta x^3 + \frac{9\alpha_2^2}{4}x^2 + 4\alpha_0\beta x + 4\beta + 2\alpha_2\alpha_0,$$

where C_1 and C_0 are arbitrary constants. If we take $C_1 = 0$ and $C_0 = 0$, then the spectral curve of operators L, M has the form

$$16w^2 - 8w(\alpha_2\alpha_0 - 2\beta + 6z^2) + 32z^4 + 16(\alpha_2\alpha_0 - 2\beta)z^2 + \alpha_2^2\alpha_0^2 = 0$$

We see that the spectral curve is nonsingular for almost all $\alpha_2, \alpha_0.\beta$ and L, M are operators of rank 2. If $\alpha_0 = 0$, then the spectral curve has the form

$$(w - 2z^2)(w - z^2 + \beta) = 0$$

and L, M are operators of vector rank $(2, 2)$.

Theorem 3. *Let $\wp(x)$ be the Weierstrass elliptic function satisfying the equation $(\wp'(x))^2 = 4\wp^3(x) + g_2\wp(x)$. The operator*

$$L = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \partial_x^2 + \begin{pmatrix} 0 & \alpha\wp(x) \\ \alpha\wp(x) & 0 \end{pmatrix},$$
$$\alpha^2 = 64n^4 - 4n^2, \quad n \in \mathbb{N}$$

commutes with a differential operator (14), of order $4n$, where $g = 2n$. The order of operator M equals $4n$.

Proofs of Theorem 2 and Theorem 3

We prove Theorem 2 and Theorem 3 using Theorem 1. Let us suppose that $C_2^k = C_3^k = C_4^k = 0$ and $C_1^{2k+1} = 0$ for all k. We know that

$$A_0 \equiv 0, \quad B_0 = \begin{pmatrix} \mu_1 & 0 \\ 0 & \mu_2 \end{pmatrix}.$$

Direct calculations using $(6)-(13)$ show that

$$\begin{gathered} a_1^1 = a_2^1 = a_3^1 = a_4^1 = 0, \\ b_2^1 = \frac{\mu_1 - \mu_2}{2} q_2, \quad b_3^1 = -\frac{\mu_1 - \mu_2}{2} q_2 = -b_2^1. \end{gathered} \tag{16}$$

Then

$$\begin{gathered} a_1^2 = a_4^2 = 0, \quad a_2^2 = a_3^2 = -\frac{\mu_1 - \mu_2}{2} q_2' = -(b_2^1)', \\ b_2^2 = b_3^2 = \frac{r_1 a_2^2 - (a_2^2)'}{2} \\ a_1^3 = a_2^3 = a_3^3 = a_4^3 = 0, \\ b_2^3 = -b_3^3 \end{gathered} \tag{17}$$

Lemma 1. If $k = 2m+1$, then

$$a_1^k = a_2^k = a_3^k = a_4^k = 0, \quad b_2^k = -b_3^k \tag{18}$$

If $k = 2m$, then

$$a_1^k = a_4^k = 0, \quad a_2^k = a_3^k = -(b_2^{k-1})', \quad b_3^k = b_2^k = \frac{r_1 a_2^k - (a_2^k)'}{2} \tag{19}$$

Proof
We see that relations (18) and (19) is true when $k = 1$ and $k = 2$. Let us suppose that (18) and (19) is true for some k.
If $k = 2m+1$, then using $(6)-(13)$ we get

$$a_1^{2m+2} = a_4^{2m+2} = 0, \quad a_2^{2m+2} = a_3^{2m+2} = -(b_2^{2m+1})', \quad b_3^{2m+2} = b_2^{2m+2} = \frac{r_1 a_2^{2m+2} - (a_2^{2m+2})'}{2}.$$

If $k = 2m$, then again using $(6)-(13)$ we have

$$a_1^{2m+1} = a_2^{2m+1} = a_3^{2m+1} = a_4^{2m+1} = 0 \qquad b_3^{2m+1} = -b_2^{2m+1}.$$

The Lemma is proved.

Proof of Theorem 2

From (16) and (17) we get

$$\begin{gathered} a_1^1 = a_2^1 = a_3^1 = a_4^1 = 0, \\ b_2^1 = \frac{\mu_1 - \mu_2}{2} q_2 = \frac{\mu_1 - \mu_2}{2} \gamma x, \quad b_3^1 = -\frac{\mu_1 - \mu_2}{2} q_2 = -\frac{\mu_1 - \mu_2}{2} \gamma x. \end{gathered}$$

Then

$$\begin{gathered} a_1^2 = a_4^2 = 0, \quad a_2^2 = a_3^2 = -\frac{\mu_1 - \mu_2}{2}\gamma = -(b_2^1)', \\ b_2^2 = b_3^2 = -\frac{\mu_1 - \mu_2}{4}\alpha_0\gamma - \frac{\mu_1 - \mu_2}{4}\alpha_2\gamma x^2, \\ a_1^3 = a_2^3 = a_3^3 = a_4^3 = 0, \\ b_2^3 = -b_3^3 = \frac{2C_1^2 + (\mu_1 - \mu_2)(\alpha_0\alpha_2 - 2\beta)}{4}\gamma x + \frac{\mu_1 - \mu_2}{4}\gamma(\gamma^2 + \alpha_2^2)x^3 \end{gathered} \tag{20}$$

We want to prove that L commutes with differential operator (14), where $g = 2n$. From Theorem 1 and Lemma 1 we know that we must prove that there exists constants C_1^{2k} such that $b_2^{2n+1} \equiv 0$. Let us note that recurrence relations $(6) - (13)$ are linear in a_i^{k+1} and b_i^{k+1}. Assume that $b_i^{2m-1} = x^{2m-1}$. Then we have

$$\begin{gathered} b_2^{2m} = (2m-1)(m-1)x^{2m-3} - \frac{\alpha_0(2m-1)x^{2m-2}}{2} - \frac{\alpha_2(2m-1)x^{2m}}{2} = -b_3^{2m}, \\ a_2^{2m} = -(b_2^{2m-1})' = -(2m-1)x^{2m-2} = a_3^{2m}. \end{gathered}$$

Again using $(6) - (13)$ we obtain

$$\begin{gathered} a_1^{2m+1} = a_2^{2m+1} = a_3^{2m+1} = a_4^{2m+1} = 0, \\ b_2^{2m+1} = \frac{(2m-1)(\alpha_2^2 m^2 + \gamma^2)}{2m}x^{2m+1} + \\ + \frac{(\alpha_2\alpha_0 - 2\beta)(2m-1)^2}{2}x^{2m-1} + \frac{\alpha_0^2(2m-2)(2m-1)}{4}x^{2m-3} - \\ - \frac{(2m-1)(2m-2)(2m-3)(2m-4)}{4}x^{2m-5} + \frac{C_1^{2m}\beta}{2}x. \end{gathered} \tag{21}$$

From (20) we see that $b_2^3 = K_1^3 x + K_3^3 x^3$, where $K_1^3 = K_1^3(C_1^2)$ is constant and depends on C_1^2, $K_3^3 = \frac{\mu_1 - \mu_2}{4}\gamma(\gamma^2 + \alpha_2^2)$. Let us suppose that for some m

$$b_2^{2m-1} = K_1^{2m-1}x + K_3^{2m-1}x^3 + ... + K_{2m-1}^{2m-1}x^{2m-1},$$

where

$$K_{2m-1}^{2m-1} = -\frac{(\mu_1 - \mu_2)\prod_{j=1}^{m-1}((2j-1)(\alpha_2 j^2 + \gamma^2))}{2^m(m-1)!},$$

K_{2m-3}^{2m-1} is constant and depends on C_1^2, K_{2m-5}^{2m-1} is constant and depends on C_1^2, C_1^4, $K_{2m-2j-1}^{2m-1}$ depends on $C_1^2, ..., C_1^{2j}$ and K_1^{2m-1} depends on $C_1^2, C_1^4, ..., C_1^{2m-2}$. We see

that it is true when $m = 2$. Using (21) we get

$$b_2^{2m+1} = K_1^{2m+1}x + K_3^{2m+1}x^3 + ... + K_{2m-1}^{2m+1}x^{2m-1} + K_{2m+1}^{2m+1}x^{2m+1},$$

$$K_1^{2m+1} = \frac{C_1^{2m}\beta}{2} - 30K_5^{2m-1} + \frac{3\alpha_0^2}{2}K_3^{2m-1} + \frac{\alpha_2\alpha_0 - 2\beta}{2}K_1^{2m-1}$$

$$K_3^{2m+1} = \frac{\alpha_2^2 + \gamma^2}{2}K_1^{2m-1} + \frac{9(\alpha_2\alpha_0 - 2\beta)}{2}K_3^{2m-1} + 5\alpha_0^2 K_5^{2m-1} - 210K_7^{2m-1}$$

...

$$K_{2m-1}^{2m+1} = \frac{(2m-3)(\alpha_2^2(m-1)^2 + \gamma^2)}{2m-2}K_{2m-3}^{2m-1} + \frac{(\alpha_2\alpha_0 - 2\beta)(2m-1)^2}{2}K_{2m-1}^{2m-1}$$

$$K_{2m+1}^{2m+1} = \frac{(2m-1)(\alpha_2^2 m^2 + \gamma^2)}{2m}K_{2m-1}^{2m-1}.$$

Easy to see that K_{2m-1}^{2m+1} depends on constant of integration C_1^2 because K_{2m-3}^{2m-1} depends on C_1^2. And K_{2m-3}^{2m+1} depends on C_1^2, C_3^4. The last coefficient K_1^{2m+1} depends on constants of integrations $C_1^2, ..., C_1^{2m}$.
Now let us consider

$$b_2^{2n+1} = K_1^{2n+1}x + K_3^{2n+1}x^3 + ... + K_{2n-1}^{2n+1}x^{2n-1} + K_{2n+1}^{2n+1}x^{2n+1}$$

We know that $\gamma^2 + n^2\alpha_2 = 0$ and hence $K_{2n+1}^{2n+1} = 0$. To prove Theorem 2 we must find constants $C_1^2, ..., C_1^{2n}$ such that $K_1^{2n+1} = K_3^{2n+1} = ... = K_{2n-1}^{2n+1} = 0$. It is always possible because K_{2n-1}^{2n+1} depends n C_1^2 and K_{2n-3}^{2n+1} depends on C_1^2, C_3^2 etc. The last coefficient K_1^{2n+1} depends on constants of integration $C_1^2, ..., C_1^{2n}$.

Theorem 2 is proved.

Proof of Theorem 3

The proof of Theorem 3 coincides with the proof of Theorem 2. Let us prove that principle parts of functions a_i^{2n+1} and b_i^{2n+1} equals zero. We see from $(6) - (13)$ that a_i^j and b_i^j are elliptic functions for any i, j. Hence if principle parts of functions a_i^{2n+1} and b_i^{2n+1} equals zero, then a_i^{2n+1} and b_i^{2n+1} havn't poles and hence are constants. In our case these constants are zeroes.

From (16) and (17) we get

$$a_1^1 = a_2^1 = a_3^1 = a_4^1 = 0,$$

$$b_2^1 = \frac{\mu_1 - \mu_2}{2}q_2 = \frac{\mu_1 - \mu_2}{2x^2}\alpha + O(x^2), \quad b_3^1 = -\frac{\mu_1 - \mu_2}{2x^2}\alpha + O(x^2).$$

Then

$$a_1^2 = a_4^2 = 0, \quad a_2^2 = a_3^2 = \frac{\mu_1 - \mu_2}{x^3}\alpha + O(x) = -(b_2^1)',$$
$$b_2^2 = b_3^2 = \frac{3(\mu_1 - \mu_2)}{2x^4}\alpha + \frac{(\mu_1 - \mu_2)g_2\alpha}{40} + O(x^4),$$
$$a_1^3 = a_2^3 = a_3^3 = a_4^3 = 0,$$
$$b_2^3 = -b_3^3 = \frac{(\mu_1 - \mu_2)\alpha(\alpha^2 - 60)}{4x^6} + \frac{\alpha(80C_1^2 + 6(\mu_1 - \mu_2)g_2\alpha^2)}{160x^2} + O(x^2) \tag{22}$$

We mentioned before that recurrence relations $(6) - (13)$ are linear in a_i^{k+1} and b_i^{k+1}. Assume that $b_3^{2m-1} = -b_2^{2m-1} = \frac{1}{x^{4m-2}}$ and $a_1^{2m-1} = a_2^{2m-1} = a_3^{2m-1} = a_4^{2m-1} = 0$. Then we have

$$b_2^{2m} = b_3^{2m} = \frac{8m^2 - 6m + 1}{x^{4m}},$$
$$a_2^{2m} = -(b_2^{2m-1})' = \frac{4m - 2}{x^{4m-1}}.$$

Again using $(6) - (13)$ we obtain

$$a_1^{2m+1} = a_2^{2m+1} = a_3^{2m+1} = a_4^{2m+1} = 0,$$
$$b_2^{2m+1} = \frac{(2m-1)(\alpha^2 + 4m^2 - 64m^4)}{2mx^{4m+2}} + \frac{const}{x^{4m-2}} + \frac{const}{x^{4m-6}} + ... + \frac{C_1^{2m}\alpha}{2x^2} + O(x^2) \tag{23}$$

From (22) we see that $b_2^3 = \frac{K_6^3}{x^6} + \frac{K_2^3}{x^2} + O(x^2)$, where $K_2^3 = K_2^3(C_1^2)$ is constant and depends on C_1^2, $K_6^3 = \frac{(\mu_1 - \mu_2)\alpha(\alpha^2 - 60)}{4}$. Let us suppose that for some m

$$b_2^{2m-1} = \frac{K_{4m-2}^{2m-1}}{x^{4m-2}} + \frac{K_{4m-6}^{2m-1}}{x^{4m-6}} + ... + \frac{K_2^{2m-1}}{x^2} + O(x^2),$$

where

$$K_{4m-2}^{2m-1} = -\frac{(\mu_1 - \mu_2)\prod_{j=1}^{m-1}((2j-1)(\alpha^2 + 4j^2 - 64j^4)))}{2^m(m-1)!},$$

K_{4m-6}^{2m-1} is constant and depends on C_1^2, K_{4m-10}^{2m-1} is constant and depends on C_1^2, C_1^4, $K_{4m-4j-2}^{2m-1}$ depends on $C_1^2, ..., C_1^{2j}$ and K_2^{2m-1} depends on $C_1^2, C_1^4, ..., C_1^{2m-2}$. We see that it is true when $m = 2$. Using (23) we get

$$b_2^{2m+1} = K_1^{2m+1}x + K_3^{2m+1}x^3 + ... + K_{2m-1}^{2m+1}x^{2m-1} + K_{2m+1}^{2m+1}x^{2m+1}.$$

Easy to see that K_{4m-2}^{2m+1} depends on constant of integration C_1^2. And K_{4m-6}^{2m+1} depends on C_1^2, C_3^2. The last coefficient K_2^{2m+1} depends on constants of integrations $C_1^2, ..., C_1^{2m}$.
Now let us consider

$$b_2^{2n+1} = K_1^{2n+1}x + K_3^{2n+1}x^3 + ... + K_{2n-1}^{2n+1}x^{2n-1} + K_{2n+1}^{2n+1}x^{2n+1}$$

We know that $\alpha^2 + 4n^2 - 64n^4 = 0$ hence $K_{2n+1}^{4n+2} = 0$. To prove Theorem 3 we must find constants $C_1^2, ..., C_1^{2n}$ such that $K_2^{2n+1} = K_6^{2n+1} = ... = K_{4n-2}^{2n+1} = 0$. It is always possible because K_{4n-2}^{2n+1} depends on C_1^2 and K_{4n-2}^{2n+1} depends on C_1^2, C_3^2 etc. The last coefficient K_2^{2n+1} depends on constants of integration $C_1^2, ..., C_1^{2n}$.

Theorem 3 is proved.

References

[1] Wallenberg, G.: Uber die Vertauschbarkeit homogener linearer Differentialausdrucke. Arch. Math. Phys. 4 (1903), 252–268.

[2] Schur, J.: Uber vertauschbare lineare Differentialausdrucke. Sitzungsber. der Berliner Math. Gesell. 4 (1905), 2–8.

[3] Burchnall J.-L., Chaundy T.W. Commutative ordinary differential operators. Proc. London Math. Soc. 21 (1923), 420-440; Proc. Royal Soc. London (A) 118 (1928), 557-583.

[4] Grinevich, P. G.: Vector rank of commuting matrix differential operators. Proof of S. P. Novikov's criterion, Math USSR IZV, 1987, 28 (3), 445–465.

[5] Krichever, I. M.: Integration of nonlinear equations by the methods of algebraic geometry, Functional Analysis and Its Applications, 11: 1 (1977), 12–26.

[6] Dixmier J.: Sur les algebres de Weyl. Bulletin de la Societe Mathematique de France 96, 209–242 (1968)

[7] Krichever, I.M.: Commutative rings of ordinary linear differential operators, Functional Functional Analysis and Its Applications, 12:3 (1978), 175–185.

[8] Krichever, I. M., Novikov, S.P.: Holomorphic bundles over algebraic curves and non-linear equations, Russian Mathematical Surveys, 1980, 35:6, 53–79.

[9] Mokhov, O. I.: Commuting ordinary differential operators of rank 3 corresponding to an elliptic curve, Russian Math. Surveys, 37:4 (1982), 129–130.

[10] Mokhov, O. I.: Commuting differential operators of rank 3, and nonlinear differential equations, Math. USSR, Izvestiya,35:3 (1990), 629–655.

[11] Mironov, A. E.: Self-adjoint commuting differential operators and commutative subalgebras of the Weyl algebra, Invent. math. (2014) 197:417-431.

[12] Mironov, A. E.: Periodic and rapid decay rank two self-adjoint commuting differential operators, Amer. Math. Soc. Transl. Ser. 2, V. 234, 2014, P. 309–322.

[13] Oganesyan, V.: Commuting differential operators of rank 2 with polynomial coefficients, Functional Analysis and Its Applications, 2016, 50:1, 54–61.

[14] Oganesyan, V.: Commuting differential operators of rank 2 and arbitrary genus g with polynomial coefficients, International Mathematics Research Notices (2016), doi:10.1093/imrn/rnw085.

[15] Davletshina V.N.: Commuting differential operators of rank 2 with trigonometric coeffcients, Siberian Mathematical Journal, 56, 405-410 (2015)

[16] Mironov, A. E., Zheglov, A. B.: Commuting Ordinary Differential Operators with Polynomial Coefficients and Automorphisms of the First Weyl Algebra, International Mathematics Research Notices 2015 : rnv218.

[17] Mokhov, O. I.: Commuting ordinary differential operators of arbitrary genus and arbitrary rank with polynomial coefficients, American Mathematical Society Translations, Volume 234 (2014), 323-336.

[18] Mokhov, O. I.: On Commutative Subalgebras of the Weyl Algebra Related to Commuting Operators of Arbitrary Rank and Genus, Mathematical Notes, vol. 94:2 (2013), 298-300.

[19] Smirnov, A. O.: Finite-gap elliptic solutions of the KdV equation. Acta Appl. Math., 36 (1994), 125-166.

[20] Smirnov, A. O.: Real finite-gap regular solutions of the Kaup-Boussinesq equation, Theoret. and Math. Phys., 66 (1986), 19-31.

[21] Dubrovin, B. A, Natanzon, S. M.: Real theta-function solutions of the Kadomtsev–Petviashvili equation, Mathematics of the USSR-Izvestiya, 1989, 32:2, 269–288.

[22] Dubrovin, B. A, Natanzon, S. M.: Real two-zone solutions of the sine-Gordon equation, Functional Analysis and Its Applications, 1982, 16:1, 21–33.

[23] Dubrovin, B. A.: The kadomcev-petviasvili equation and the relations between the periods of holomorphic differentials on riemann surfaces, Mathematics of the USSR-Izvestiya, 1982, 19:2, 285–296.

[24] Shiota, T., Characterization of Jacobian varieties in terms of soliton equations. Invent. Math. 83 (1986), no. 2, 333–382.

[25] A. Belov-Kanel, M. Kontsevich, The Jacobian conjecture is stably equivalent to the Dixmier conjecture, Moscow Mathematical Journal 7 (2): 209–218.

[26] Y. Tsuchimoto, Endomorphisms of Weyl algebra and p-curvatures, Osaka J. Math. 42: 435–452.

[27] Gesztesy, F., Weikard, R.: A characterization of all elliptic algebro-geometric solutions of the AKNS hierarchy, Acta Mathematica, September 1998, Volume 181, Issue 1, pp 63-108.

[28] Weikard, R.: On commuting matrix differential operators, New York J. Math., vol. 8, 2002, pp. 9-30.

[29] Dubrovin, B. A.: Completely integrable Hamiltonian systems associated with matrix operators and Abelian varieties, Functional Analysis and Its Applications, 1977, 11:4, 265–277.

[30] Krichever, I.M.: Algebraic curves and commuting matricial differential operators, Functional Analysis and Its Applications, 1976, 10:2, 144–146.

Department of Geometry and Topology, Faculty of Mechanics and Mathematics, Lomonosov Moscow State University, Moscow, 119991 Russia.

E-mail address: vardan.o@mail.ru